PRACTICAL CLINICAL COURSES

A Service of the Gordon J. Christensen Career Development Program

V3105 Uncomplicated Occlusal Equilibration

Gordon J. Christensen, DDS, MSD, PhD

Materials Included:

Terminology Confusion Clarification
C.E. Instruction Sheet
Products List
Clinician Responsible
Goals & Objectives
Overview
Supplemental Materials
AGD Post-Test

TERMINOLOGY CONFUSION CLARIFICATION

The glossary of Prosthodontic terms defines CENTRIC OCCLUSION as the "occlusion of opposing teeth when the mandible is in centric relation. This may or may not coincide with maximum intercuspal position."

The glossary definition is not the one most dentists have learned, and it may be confusing on the video you are viewing. Previous definitions of CENTRIC OCCLUSION have indicated that this position is the location where the patient chews, regardless of where it is in regard to centric relation.

In this video please interpret the phrase CENTRIC OCCLUSION, which is the older phrase that most dentists use, to mean the MAXIMAL INTERCUSPAL POSITION or MIP, which is the best fit of the teeth regardless of the condylar position.

It is my plan to eventually eliminate the phrase centric occlusion and replace it with the phrase maximal intercuspal position or MI.

Sorry for the confusion!

Thank you!

Gordon Christensen

Gordon J. Christensen PRACTICAL CLINICAL COURSES

PROCEDURE FOR RECEIVING CE CREDIT FOR VIDEO COURSES

- 1. Complete the enclosed Post-Test.* For each <u>CE Video Purchased</u>, one complimentary test is included. Additional tests are \$35/person per test. Fees can be paid by check or with a credit card when submitting tests to Practical Clinical Courses. If more than 10 CE tests are required per video, please contact us at 800-223-6569.
- 2. Complete the demographic information located at the end of the test.

Type of Credit:

- a. If the applicant selects "State," PCC will issue a certificate of verification to the applicant. The applicant must then submit this certificate to their state board to obtain credit.
- b. If the applicant selects "AGD," PCC will submit credit information to the Academy of General Dentistry and confirmation to the applicant that it has been submitted (the applicant may check their AGD transcript for verification please allow 30 days).
- c. If the applicant selects "Both," PCC will complete a. & b. above.
- 3. Return the **Post-Test portion** via mail, fax, or email. Our contact information is as follows:

Practical Clinical Courses 3707 N Canyon Road Suite 3D Provo, UT 84604 Fax: (801) 226-8637 cetests@pccdental.com

4. Practical Clinical Courses will correct the Post-Test. Passing scores are 70% or higher.

*TO OBTAIN CE CREDIT ONLINE: Login or create an account on www.pccdental.com and select "My CE Tests" from the left-side menu. Click on the video title to take the test online. RESULTS ARE IMMEDIATE. Missing the test? Contact us at 800-223-6569 during our business hours of 7:00 a.m. – 4:00 p.m. MST to add it to your account.

Gordon J. Christensen

PRACTICAL CLINICAL COURSES

Sources of Products Discussed in

V3105 Uncomplicated Occlusal Equilibration

Presented by: Gordon J. Christensen, DDS, MSD, PhD

1. ACCUFILM II

Parkell 155 Schmitt Blvd. P.O. Box 376 Farmingdale, NY 11735 (800)243-7446 (631)249-1134 www.parkell.com

2. ARTICULATING PAPER FORCEPS

Local Retailer

3. ARTICULATOR 2240Q

Whip Mix Corporation 361 Farmington Avenue P.O. Box 17183 Louisville, KY 40217 (800)626-5651 (502)637-1451 www.whipmix.com

4. 7406 TRIMMING & FINISHING BUR

Dentsply Professional 1301 Smile Way P.O. Box 7807 York, PA 17404-0807 (800)989-8826 (717)767-8500 www.dentsply.com

5. COTTON ROLLS (6 INCH AND REGULAR)

Local Retailer

6. DRY-TIPS

Local Retailer

7. MADAME BUTTERFLY SILK

Almore International, Inc. P.O. Box 25214 Portland, OR 97298-0214 (800)547-1511 (503)643-6633 www.almore.com

8. OCCLUSAL INDICATOR WAX

Kerr Corporation 1717 West Collins Orange, CA 92867 (800)537-7123 (714)516-7400 www.kerrdental.com

9. PRISMA GLOSS

Dentsply Caulk 38 West Clarke Avenue P.O. Box 359 Milford, DE 19963-0359 (800)532-2855 (302)422-4511 www.caulk.com

10. PROPHY PASTE

Dentsply Professional 1301 Smile Way P.O. Box 7807 York, PA 17404-0807 (800)989-8826 (717)767-8500 www.dentsply.com

Product names, the products themselves, and company names change rapidly. Please contact the companies shown to confirm current information.

Gordon J. Christensen Practical Clinical Courses, 3707 North Canyon Road, Suite 3D, Provo, UT 84604 Toll Free (800) 223-6569 or Utah Residents (801) 226-6569

The techniques and procedures on this videotape are intended to be suggestions only. Any licensed practitioner viewing this presentation must make his or her own professional decisions about specific treatment for patients. PCC is not responsible for any damages or other liabilities (including attorney's fees) resulting, or claimed to result in whole or in part, from actual or alleged problems arising out of the use of this presentation.

PROGRAM

V3105 Uncomplicated Occlusal Equilibration

CLINICIAN RESPONSIBLE:

Gordon J. Christensen, DDS, MSD, PhD

Founder and CEO, Practical Clinical Courses Senior Consultant & Previous CEO, CR Foundation Practicing Prosthodontist, Provo, Utah

GOALS & OBJECTIVES

At the completion of this video presentation, participants should be able to accomplish the following:

- 1. Explain the importance of occlusal equilibration.
- 2. Describe normal occlusion in a young adult.
- 3. Define centric occlusion.
- 4. Define centric relation.
- 5. Define working.
- 6. Define non-working.
- 7. Define long-centric.
- 8. Define wide-centric.
- 9. Relate incisal guidance to protrusive.
- 10. List five indications for occlusal equilibration.
- 11. Discuss the problems that occur if occlusal equilibration is not accomplished for the five indications you listed in statement 10.
- 12. List the instruments and materials necessary for occlusal equilibration.
- 13. Describe the difference in tooth roughness/smoothness between use of a diamond or 12-bladed cutting instrument.
- 14. Describe the clinical steps in the occlusal equilibration technique.
- 15. Describe why the patient is adjusted in a supine position and also in a sitting-up position.
- 16. Describe the most common location for initial contact of teeth in centric relation.
- 17. Describe why cusps are usually not cut in an occlusal equilibration.
- 18. Describe why marginal ridges, transverse ridges, and central fossa are reduced when accomplishing an occlusal equilibration.
- 19. Discuss treatment if tooth sensitivity occurs after an occlusal equilibration.
- 20. Describe the difference between a CRO occlusal equilibration and a long-centric/wide-centric occlusal equilibration.

OVERVIEW

V3105 Uncomplicated Occlusal Equilibration

Most dentists have had an introduction to occlusal equilibration while in dental school. However, it was taught in the onrush of clinical procedures that were required for graduation, with the result being that many practitioners do not have a practical approach to occlusal equilibration. This presentation emphasizes a simplified concept of occlusal equilibration that can be implemented into practice easily and accomplished frequently.

Occlusal equilibration is necessary as a part of the overall therapy in the following treatment situations: placement of routine tooth restorations (operative dentistry); orthodontics; periodontics; implant dentistry; fixed and removable prosthodontics; endodontics; and other areas. If partial or complete occlusal equilibration is not accomplished as a part of the preceding treatments, various maladies occur ranging from broken teeth, to the necessity for endodontic therapy, to relapse of orthodontic therapy, or rejection of implants.

It is necessary to know the terminology of occlusion to communicate with other practitioners. The following terms are described and demonstrated in the presentation: centric occlusion, centric relation, working, non-working, protrusive, wide-centric, long-centric, incisal guidance, group function, and canine rise.

The presentation emphasizes and demonstrates establishment of a centric-relation-occlusion (CRO) chewing position during the occlusal equilibration treatment. Long/wide centric occlusal equilibrations are also described and their indications noted. The following indications for an occlusal equilibration are described: pre-restorative; during and post-restorative, as a part of periodontal therapy, as a part of orthodontic therapy, as a part of temporomandibular joint therapy, and for bruxism and clenching. The following steps are suggested and shown for a (CRO) occlusal equilibration:

- 1. Adjust centric relation with patient in a supine position.
- 2. Patient rest for a few minutes while sitting up.
- 3. Adjust centric relation with patient in a sitting position.
- 4. Adjust working position with patient in a supine position.
- 5. Adjust non-working position with patient in a supine position.
- 6. Adjust protrusive with patient in a supine position.
- 7. Patient rest for a few minutes while sitting up.
- 8. Check and correct all positions while patient sits up.
- 9. Reappoint patient for occlusal "touch-up" about 10 days in the future.
- 10. Reappoint patient for a second occlusal "touch-up" 20 days in the future.

Occlusal equilibration is a necessary part of many dental therapies, and it is neglected in many situations.

This presentation prepares practitioners to accomplish simple occlusal equilibrations with ease.

SUPPLEMENTAL MATERIALS

V3105 Uncomplicated Occlusal Equilibration

- 1. CHRISTENSEN, G.J. "Abnormal Occlusal Conditions: A Forgotten Part of Dentistry". <u>JADA</u>. Vol. 126, December 1995. Pp. 1667-1668.
- 2. CHRISTENSEN, G.J. "Treating Bruxism and Clenching". JADA. Vol. 131, February 2000. Pp. 233-235.
- 3. CHRISTENSEN, G.J. "Now is the Time to Observe and Treat Dental Occlusion". **JADA**. Vol. 132, January 2001. Pp. 100-102.
- 4. CHRISTENSEN, G.J. "Is Occlusion Becoming More Confusing? A Plea for Simplicity". <u>JADA</u>. Vol. 135, June 2004. Pp. 767-770.

POST-TEST

V3105 Uncomplicated Occlusal Equilibration

- 1. Mark the one *incorrect statement*:
 - a. Centric occlusion is the most intercuspated position.
 - b. Centric relation is one mm anterior to centric occlusion.
 - c. Incisal guidance is present in almost all patients.
 - d. Working occurs on the chewing side.
 - e. Non-working is on the non-chewing side.
- 2. Non-working contacts with opposing teeth:
 - a. should be left intact in the natural dentition.
 - b. should be removed in the natural dentition.
 - c. working contacts should be removed in the natural dentition.
 - d. when working contacts are reduced, non-working contacts are decreased also.
- 3. Protrusive contacts are:
 - a. almost always present in the natural dentition.
 - b. should be reduced on most patients.
 - c. are usually on the anterior and premolar teeth only.
 - d. usually not homogeneous across the anterior segment.
- 4. A shift forward from centric relation to centric occlusion is:
 - a. always present.
 - b. about 3 mm forward on the average.
 - c. always pathogenic.
 - d. about 1 mm forward and upward on the average.
- 5. Casts mounted on a semi-adjustable articulator should:
 - a. be used for study each time before an occlusal equilibration.
 - b. be considered as a good learning experience, but are usually not necessary when accomplishing an occlusal equilibration.
 - c. be mounted in centric occlusion.
 - d. be mounted in protrusive.
- 6. Select the *incorrect statement*. Occlusal equilibration:
 - a. should start with centric relation adjustment.
 - b. should be accomplished partly in the supine position and partly with the patient sitting up.
 - c. should be considered to be finished and permanent after the second touch-up appointment.
 - d. should be explained carefully to patients before starting the procedure.
- 7. Select the *incorrect statement*. Cutting and smoothing teeth should be accomplished with:
 - a. a 7406 twelve-bladed bur.
 - b. Prophy Paste.
 - c. composite polishing paste.
 - d. a medium grit diamond rotary instrument.

POST-TEST (CONT'D)

V3105 Uncomplicated Occlusal Equilibration

- 8. Bruxism and clenching:
 - a. can be stopped by occlusal equilibration.
 - b. are the same conditions.
 - c. can be slowed by occlusal equilibration.
 - d. are not indications for occlusal equilibration.
- 9. Moving the mandible into centric relation position was suggested to be done by:
 - a. dual-handed mandibular manipulation.
 - b. single-handed mandibular manipulation.
 - c. an electronic device.
 - d. both the dental assistant and the dentist manipulating the jaw at the same time.
- 10. If it appears that accomplishing a CRO occlusal equilibration will cause cutting through crowns or enamel:
 - a. discontinue the equilibration.
 - b. adjust only non-working.
 - c. always tell the patient new crowns are needed.
 - d. accomplish a partial occlusal adjustment, with a remaining slide from centric relation to centric occlusion.

PLEASE PRINT

Name	
Name of video purchaser (if different from above)	
Address	
City/State/Zip	
Phone No	
Email	
Indicate which type of credit you wish to obtain: □ AGD □ State □ Both	
State License No AGD No	
Payment information Visa American Express Mastercard Discover	
Card # Expires / CVV2-Code:	

For each <u>CE Video Purchased</u>, one complimentary test is included. If you require CE tests for *staff members or additional doctors to receive credit, the fee is \$35/person per test*.

If more than 10 CE tests are required per video, please contact PCC at 800-223-6569.